



**Kanuma**<sup>™</sup>  
sebelipase alfa  
intravenous infusion

FOR TREATMENT OF PATIENTS WITH LYSOSOMAL ACID LIPASE  
DEFICIENCY (LAL-D)

# **KANUMA**<sup>™</sup> (sebelipase alfa) **DOSING AND ADMINISTRATION GUIDE**



**Please see Important Safety Information  
on page 9 and the attached full  
Prescribing Information for KANUMA.**

## INDICATION



**KANUMA™ (sebelipase alfa) is indicated for the treatment of patients with a diagnosis of LAL-D.**

## RECOMMENDED DOSE OF KANUMA

The recommended dose of KANUMA is based on patient weight and rate of progression of LAL-D.



For **Infants (under 6 months of age) with rapidly progressive LAL-D**, the recommended starting dosage is 1 mg/kg administered once weekly as an intravenous infusion. For patients who do not achieve an optimal clinical response, increase dosage to **3 mg/kg** once weekly.



For **Pediatric and Adult patients with LAL-D**, the recommended dosage is **1 mg/kg administered once every other week** as an intravenous infusion.

### IMPORTANT SAFETY INFORMATION

#### WARNINGS AND PRECAUTIONS

##### Hypersensitivity Reactions Including Anaphylaxis:

Hypersensitivity reactions, including anaphylaxis, have been reported in KANUMA-treated patients. In clinical trials, 3 of 106 (3%) patients treated with KANUMA experienced signs and symptoms consistent with anaphylaxis. These patients experienced reactions during infusion with signs and symptoms including chest discomfort, conjunctival injection, dyspnea, generalized and itchy rash, hyperemia, swelling of eyelids, rhinorrhea, severe respiratory distress, tachycardia, tachypnea, and urticaria. Anaphylaxis has occurred as early as the sixth infusion and as late as 1 year after treatment initiation.

**Please see Important Safety Information on page 9 and the attached full Prescribing Information for KANUMA.**

## CALCULATING THE AMOUNT OF KANUMA NEEDED

KANUMA is a concentrated solution for intravenous infusion after dilution; KANUMA comes in a 10-mL, single-use vial containing 20 mg of sebelipase alfa.

Determine the number of vials to be diluted based on the patient's weight and the recommended dose.

### 1. TO CALCULATE THE TOTAL VOLUME OF KANUMA THE PATIENT NEEDS:

#### For patients receiving a dose of 1 mg/kg:

$$\text{Patient weight (kg)} \times 1 \text{ mg/kg dose} \times 1 \text{ mL/2 mg} = \text{Total mL of KANUMA}$$

#### For infants with rapidly progressive LAL-D receiving an escalated dose of 3 mg/kg:

$$\text{Patient weight (kg)} \times 3 \text{ mg/kg dose} \times 1 \text{ mL/2 mg} = \text{Total mL of KANUMA}$$

### 2. TO CALCULATE THE NUMBER OF VIALS OF KANUMA THE PATIENT NEEDS:

$$\text{Total mL of KANUMA} \div 10 \text{ mL per vial} = \text{Total number of vials (round up to the next whole vial)}$$

Note: Convert a patient weight in pounds (lb) to kilograms (kg) by dividing by 2.2.

#### ▶ DOSE CALCULATION EXAMPLE FOR A PATIENT WHO WEIGHS 3 KG

$$\begin{aligned} \text{Patient weight (3 kg)} \times \text{dose (1 mg/kg)} \times 1 \text{ mL/2 mg} &= (1.5 \text{ mL of KANUMA}) \\ \text{Total volume of KANUMA (1.5 mL)} \div 10 \text{ mL per vial} &= 0.15 = 1 \text{ vial per infusion}^a \end{aligned}$$

#### ▶ EXAMPLE OF DOSE ESCALATION TO 3 MG/KG FOR A PATIENT WHO WEIGHS 3 KG

$$\begin{aligned} \text{Patient weight (3 kg)} \times \text{dose (3 mg/kg)} \times 1 \text{ mL/2 mg} &= (4.5 \text{ mL of KANUMA}) \\ \text{Total volume of KANUMA (4.5 mL)} \div 10 \text{ mL per vial} &= 0.45 = 1 \text{ vial per infusion}^a \end{aligned}$$

#### ▶ DOSE CALCULATION EXAMPLE FOR A PATIENT WHO WEIGHS 50 KG

$$\begin{aligned} \text{Patient weight (50 kg)} \times \text{dose (1 mg/kg)} \times 1 \text{ mL/2 mg} &= (25 \text{ mL of KANUMA}) \\ \text{Total volume of KANUMA (25 mL)} \div 10 \text{ mL per vial} &= 3 \text{ vials per infusion}^a \end{aligned}$$

<sup>a</sup>Discard any remaining unused KANUMA left in the vial.

## SUPPLIES NEEDED FOR INFUSION

Use the following supplies to administer KANUMA.

- A** KANUMA 10-mL, single-use vials<sup>a</sup>
- B** 0.9% Sodium Chloride Injection, USP
- C** Low-protein binding infusion set equipped with an in-line, low-protein binding 0.2 micron filter
- D** Syringe
- E** Needles (do not use filter needles)



<sup>a</sup>Contain 20 mg of sebelipase alfa.

### IMPORTANT SAFETY INFORMATION

#### Hypersensitivity Reactions Including Anaphylaxis (cont'd):

In clinical trials, 21 of 106 (20%) KANUMA-treated patients, including 9 of 14 (64%) infants and 12 of 92 (13%) pediatric patients, 4 years and older, and adults experienced signs and symptoms either consistent with or that may be related to a hypersensitivity reaction.

**Please see Important Safety Information on page 9 and the attached full Prescribing Information for KANUMA.**

## STORING KANUMA

- Vials are for single-use only
- KANUMA should be stored under refrigeration between 2°C and 8°C (36°F and 46°F)
- Do not freeze or shake
- Protect from light
- KANUMA contains no preservatives; therefore, product should be used immediately after dilution; if immediate use is not possible, the diluted product may be stored up to 24 hours under refrigeration at 2°C to 8°C (36°F to 46°F)
- Discard any unused product

## DILUTION PRIOR TO ADMINISTRATION

Prior to administration, prepare KANUMA for dilution using aseptic techniques.

KANUMA must be diluted with 0.9% Sodium Chloride Injection, USP. Use the table below to determine a patient's total infusion volume.

PATIENT WEIGHT RANGE (kg)	1 mg/kg dose	3 mg/kg dose <sup>b</sup>
	TOTAL INFUSION VOLUME (mL) <sup>c</sup>	TOTAL INFUSION VOLUME (mL) <sup>c</sup>
1-10.9	10	25
11-24.9	25	50
25-49.9	50	100
50-99.9	100	250
100-120.9	250	500

<sup>b</sup> For patients with LAL-D presenting within the first 6 months of life who do not achieve an optimal clinical response with a dose of 1 mg/kg.

<sup>c</sup>The infusion volume should be based on the prescribed dose and should be prepared to a final KANUMA concentration of 0.1 mg/kg to 1.5 mg/mL.

### IMPORTANT SAFETY INFORMATION

#### Hypersensitivity Reactions Including Anaphylaxis (cont'd):

Signs and symptoms of hypersensitivity reactions, occurring in two or more patients, included abdominal pain, agitation, fever, chills, diarrhea, eczema, edema, hypertension, irritability, laryngeal edema, nausea, pallor, pruritus, rash, and vomiting. The majority of reactions occurred during or within 4 hours of the completion of the infusion. Patients were not routinely pre-medicated prior to infusion of KANUMA in these clinical trials.



# ADMINISTER KANUMA USING THE FOLLOWING STEPS

KANUMA is administered by a healthcare professional.

- ▶ **COMPLETE THE DOSE CALCULATION** outlined on page 3 of this brochure to determine how many vials of KANUMA you will need
- ▶ **REMOVE** the calculated number of vials from the refrigerator; do not shake, heat, or freeze vials
- ▶ **SET VIALS ASIDE** to allow them to come to room temperature before diluting



**STEP 1** **MIX THE KANUMA SOLUTION GENTLY BY INVERSION** to ensure proper distribution of KANUMA. Do not shake the vials or the prepared infusion.



**STEP 2** **INSPECT EACH VIAL** visually for particulate matter and discoloration prior to administration. The solution should be a clear to slightly opalescent, colorless to slightly colored solution. Thin, translucent particles or fibers may be present in the vials or diluted solution. Do not use if the solution is cloudy or if other particulate matter is observed.

## IMPORTANT SAFETY INFORMATION

### Hypersensitivity Reactions Including Anaphylaxis (cont'd):

Due to the potential for anaphylaxis, appropriate medical support should be readily available when KANUMA is administered.

**Hypersensitivity to Eggs or Egg Products:** Consider the risks and benefits of treatment in patients with known systemic hypersensitivity reactions to eggs or egg products.

**Please see Important Safety Information on page 9 and the attached full Prescribing Information for KANUMA.**



**STEP 3** **WITHDRAW AND DISCARD** a volume of the 0.9% Sodium Chloride Injection, USP, from the infusion bag equal to the volume of KANUMA to be added. Withdraw the calculated volume of KANUMA from the corresponding number of vials and add to the infusion bag using a sterile syringe.



**STEP 4** **ADMINISTER THE DILUTED KANUMA** to the patient using a low-protein binding infusion set equipped with an in-line, low-protein binding 0.2 micron filter.

## INFUSION TIME

- ▶ **ADMINISTER THE TOTAL VOLUME OF THE INFUSION OVER AT LEAST 2 HOURS**

Consider further prolonging the infusion time for patients receiving a dose of 3 mg/kg. A 1-hour infusion may be considered for patients receiving a dose of 1 mg/kg who tolerate the infusion.

## IMPORTANT SAFETY INFORMATION

### ADVERSE REACTIONS

The most common adverse reactions are:

In patients with Rapidly Progressive Disease Presenting within the First 6 Months of Life ( $\geq 30\%$ ): diarrhea, vomiting, fever, rhinitis, anemia, cough, nasopharyngitis, and urticaria.

In pediatric and adult patients ( $\geq 8\%$ ): headache, fever, oropharyngeal pain, nasopharyngitis, asthenia, constipation, and nausea.



# SUPPORT FOR PHYSICIANS

► **LAL-D PATIENT REGISTRY:** For more information, visit [www.laldeficiencyregistry.com](http://www.laldeficiencyregistry.com) or email [help@laldeficiencyregistry.com](mailto:help@laldeficiencyregistry.com)

► **TO REPORT SUSPECTED ADVERSE REACTIONS,** contact Alexion at **1-888-765-4747** or the US Food and Drug Administration (FDA) at **1-800-332-1088** or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)

Life-threatening and severe allergic reactions may occur in patients who receive KANUMA. These reactions may occur in patients who are starting treatment with KANUMA or in patients who have previously received KANUMA without having an allergic reaction. Symptoms include:

- Chest pain or discomfort
- Wheezing or trouble breathing
- Rash or hives
- Red eyes
- Swelling of eyelids
- Rapid heartbeat
- Rapid breathing
- Runny nose

In infants with rapidly progressive disease presenting within the first 6 months of life treated with KANUMA, the most common side effects are diarrhea, vomiting, fever, stuffy or runny nose, low hemoglobin (red blood cells), cough, swelling of the nose and throat, and hives.

In pediatric and adult patients treated with KANUMA, the most common side effects are headache, fever, sore throat, swelling of the nose and throat, weakness, constipation, and nausea.

Please see Important Safety Information on page 9 and the attached full Prescribing Information for KANUMA.

## INDICATIONS AND USAGE

KANUMA™ (sebelipase alfa) is indicated for the treatment of patients with a diagnosis of LAL-D.

## IMPORTANT SAFETY INFORMATION

### CONTRAINDICATIONS

None.

### WARNINGS AND PRECAUTIONS

#### Hypersensitivity Reactions Including Anaphylaxis:

Hypersensitivity reactions, including anaphylaxis, have been reported in KANUMA-treated patients. In clinical trials, 3 of 106 (3%) patients treated with KANUMA experienced signs and symptoms consistent with anaphylaxis. These patients experienced reactions during infusion with signs and symptoms including chest discomfort, conjunctival injection, dyspnea, generalized and itchy rash, hyperemia, swelling of eyelids, rhinorrhea, severe respiratory distress, tachycardia, tachypnea, and urticaria. Anaphylaxis has occurred as early as the sixth infusion and as late as 1 year after treatment initiation.

In clinical trials, 21 of 106 (20%) KANUMA-treated patients, including 9 of 14 (64%) infants and 12 of 92 (13%) pediatric patients, 4 years and older, and adults experienced signs and symptoms either consistent with or that may be related to a hypersensitivity reaction. Signs and symptoms of hypersensitivity reactions, occurring in two or more patients, included abdominal pain, agitation, fever, chills, diarrhea, eczema, edema, hypertension, irritability, laryngeal edema, nausea, pallor, pruritus, rash, and vomiting. The majority of reactions occurred during or within 4 hours of the completion of the infusion. Patients were not routinely pre-medicated prior to infusion of KANUMA in these clinical trials.

Due to the potential for anaphylaxis, appropriate medical support should be readily available when KANUMA is administered.

**Hypersensitivity to Eggs or Egg Products:** Consider the risks and benefits of treatment in patients with known systemic hypersensitivity reactions to eggs or egg products.

### ADVERSE REACTIONS

The most common adverse reactions are:

In patients with Rapidly Progressive Disease Presenting within the First 6 Months of Life ( $\geq 30\%$ ): diarrhea, vomiting, fever, rhinitis, anemia, cough, nasopharyngitis, and urticaria.

In pediatric and adult patients ( $\geq 8\%$ ): headache, fever, oropharyngeal pain, nasopharyngitis, asthenia, constipation, and nausea.

# OneSource™ PATIENT SUPPORT PROGRAM

Once you have decided that KANUMA is right for your patient, OneSource is here to help. Alexion's objective is that every patient with LAL-D who can benefit from KANUMA will have access to it. After your patients sign up, they will be matched with a dedicated nurse case manager who can help answer any questions about KANUMA or LAL-D.

OneSource is staffed by Alexion nurse case managers who provide one-to-one education, information on funding options and access to KANUMA, and ongoing treatment support for your patients living with LAL-D, and their caregivers.

Our nurse case managers provide personalized support for your patients and their caregivers:

- Assistance from a dedicated nurse case manager
- Answers to questions about LAL-D or KANUMA
- Resources and education on the infusion process
- Help connecting patients with other families affected by LAL-D



Help get your patients connected with their OneSource nurse case manager by calling 1-888-765-4747, Monday through Friday, 8:30 AM to 5 PM (ET).

[kanuma.com](http://kanuma.com)

**Please see Important Safety Information on page 9 and the attached full Prescribing Information for KANUMA.**

**Reference:** KANUMA [package insert]. Cheshire, CT: Alexion Pharmaceuticals Inc; 2015.

**ALEXION**®

**ONESOURCE**™  
TREATMENT SUPPORT

KANUMA and OneSource are trademarks and Alexion is a registered trademark of Alexion Pharmaceuticals, Inc.  
Copyright © 2015, Alexion Pharmaceuticals, Inc.  
All rights reserved. US/KAN-LAL/15/0013

**Kanuma**™  
sebelipase alfa  
intravenous infusion

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use KANUMA safely and effectively. See full prescribing information for KANUMA.

**KANUMA (sebelipase alfa) injection, for intravenous use**  
Initial U.S. Approval: 2015

### INDICATIONS AND USAGE

KANUMA™ is a hydrolytic lysosomal cholesteryl ester and triacylglycerol-specific enzyme indicated for the treatment of patients with a diagnosis of Lysosomal Acid Lipase (LAL) deficiency. (1)

### DOSAGE AND ADMINISTRATION

**Patients with Rapidly Progressive LAL Deficiency Presenting within the First 6 Months of Life:** The recommended starting dosage is 1 mg/kg as an intravenous infusion once weekly. For patients who do not achieve an optimal clinical response, increase to 3 mg/kg once weekly. (2.1)

**Pediatric and Adult Patients with LAL Deficiency:** The recommended dosage is 1 mg/kg as an intravenous infusion once every other week. (2.1)

#### Administration Instructions (2.3):

- Infuse over at least 2 hours.
- Consider further prolonging the infusion time for the 3 mg/kg dose or if a hypersensitivity reaction occurs.
- Consider a 1-hour infusion for the 1 mg/kg dose in patients who tolerate the infusion.

### DOSAGE FORMS AND STRENGTHS

Injection: 20 mg/10 mL (2 mg/mL) solution in single-use vials. (3)

### CONTRAINDICATIONS

None. (4)

### WARNINGS AND PRECAUTIONS

- **Hypersensitivity Reactions including Anaphylaxis:** Observe patients during and after the infusion. Consider interrupting the infusion or lowering the infusion rate, based on the severity of the reaction. If a severe hypersensitivity reaction occurs, immediately stop the infusion and initiate appropriate treatment. Pre-treatment with antipyretics and/or antihistamines may prevent subsequent reactions in those cases where symptomatic treatment is required. (5.1)
- **Hypersensitivity to Eggs or Egg Products:** Consider the risks and benefits of treatment in patients with known systemic hypersensitivity reactions to eggs or egg products. (5.2)

### ADVERSE REACTIONS

The most common adverse reactions are:

- **Patients with Rapidly Progressive Disease Presenting within the First 6 Months of Life** (≥30%): diarrhea, vomiting, fever, rhinitis, anemia, cough, nasopharyngitis, and urticaria. (6.1)
- **Pediatric and Adult Patients** (≥8%): headache, fever, oropharyngeal pain, nasopharyngitis, asthenia, constipation, and nausea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Alexion at 1-844-259-6783 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 12/2015

## FULL PRESCRIBING INFORMATION: CONTENTS\*

<b>1</b>	<b>INDICATIONS AND USAGE</b>
<b>2</b>	<b>DOSAGE AND ADMINISTRATION</b>
2.1	Dosage
2.2	Preparation Instructions
2.3	Administration Instructions
2.4	Storage of Diluted Solution
<b>3</b>	<b>DOSAGE FORMS AND STRENGTHS</b>
<b>4</b>	<b>CONTRAINDICATIONS</b>
<b>5</b>	<b>WARNINGS AND PRECAUTIONS</b>
5.1	Hypersensitivity Reactions Including Anaphylaxis
5.2	Hypersensitivity to Eggs or Egg Products
<b>6</b>	<b>ADVERSE REACTIONS</b>
6.1	Clinical Trials Experience
6.2	Immunogenicity
<b>8</b>	<b>USE IN SPECIFIC POPULATIONS</b>
8.1	Pregnancy
8.2	Lactation
8.4	Pediatric Use
8.5	Geriatric Use
<b>11</b>	<b>DESCRIPTION</b>
<b>12</b>	<b>CLINICAL PHARMACOLOGY</b>
12.1	Mechanism of Action
12.2	Pharmacodynamics
12.3	Pharmacokinetics
<b>13</b>	<b>NONCLINICAL TOXICOLOGY</b>
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
13.2	Animal Toxicology and/or Pharmacology
<b>14</b>	<b>CLINICAL STUDIES</b>
14.1	Patients with Rapidly Progressive LAL Deficiency Presenting within the First 6 Months of Life
14.2	Pediatric and Adult Patients with LAL Deficiency
<b>16</b>	<b>HOW SUPPLIED/STORAGE AND HANDLING</b>
<b>17</b>	<b>PATIENT COUNSELING INFORMATION</b>

\*Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

KANUMA™ is indicated for the treatment of patients with a diagnosis of Lysosomal Acid Lipase (LAL) deficiency.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Dosage

**Patients with Rapidly Progressive LAL Deficiency Presenting within the First 6 Months of Life:**

The recommended starting dosage is 1 mg/kg administered once weekly as an intravenous infusion. For patients who do not achieve an optimal clinical response, increase to 3 mg/kg once weekly. **Pediatric and Adult Patients with LAL Deficiency:**

The recommended dosage is 1 mg/kg administered once every other week as an intravenous infusion.

#### 2.2 Preparation Instructions

KANUMA is for intravenous infusion only. Prepare KANUMA using the following steps.

1. Determine the number of vials needed based on the patient's weight and the recommended dose of 1 mg/kg or 3 mg/kg, using the following calculations (a-b):
  - a. Total dose (mg) = Patient's weight (kg) x Recommended dose (mg/kg)
  - b. Total number of vials = Total dose (mg) divided by 20 mg/vial
2. Round to the next whole vial and remove the required number of vials from the refrigerator to allow them to reach room temperature.
  - a. Volume (mL) of calculated total dose = Total dose (mg) divided by the 2 mg/mL concentration
  - b. Volume (mL) of 0.9% Sodium Chloride for dilution = Total infusion volume (mL) for patient's weight (see Table 1) - Volume (mL) of calculated total dose

Table 1: Total Infusion Volumes\*

Weight Range (kg)	1 mg/kg dose	3 mg/kg dose**
	Total Infusion Volume (mL)	Total Infusion Volume (mL)
1 to 10.9	10	25
11 to 24.9	25	50
25 to 49.9	50	100
50 to 99.9	100	250
100 to 120.9	250	500

\* The infusion volume should be based on the prescribed dose and should be prepared to a final KANUMA concentration of 0.1 mg/mL to 1.5 mg/mL.

\*\* For patients with LAL deficiency presenting within the first 6 months of life who do not achieve an optimal clinical response with a dose of 1 mg/kg.

3. Mix gently by inversion. Do not shake the vials or the prepared infusion.
4. The solution should be inspected visually for particulate matter and discoloration prior to administration. The solution should be a clear to slightly opalescent, colorless to slightly colored solution. Thin, translucent particles or fibers may be present in the vials or diluted solution. Do not use if the solution is cloudy or if other particulate matter is observed.
5. Vials are for single-use only. Discard any unused product. Do not freeze.

#### 2.3 Administration Instructions

Administer the diluted solution as an intravenous infusion using a low-protein binding infusion set with an in-line, low-protein binding 0.2 micron filter.

Infuse over at least 2 hours. Consider further prolonging the infusion time for patients receiving the 3 mg/kg dose or those who have experienced hypersensitivity reactions [see *Warnings and Precautions* (5.1)]. A 1-hour infusion may be considered for those patients receiving the 1 mg/kg dose who tolerate the infusion.

#### 2.4 Storage of Diluted Solution

KANUMA contains no preservatives; therefore, product should be used immediately after dilution. If immediate use is not possible, the diluted product may be stored up to 24 hours in the refrigerator at 2°C to 8°C (36°F to 46°F). Do not freeze or shake. Protect from light.

### 3 DOSAGE FORMS AND STRENGTHS

Injection: 20 mg/10 mL (2 mg/mL) solution in single-use vials.

### 4 CONTRAINDICATIONS

None.

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Hypersensitivity Reactions Including Anaphylaxis

Hypersensitivity reactions, including anaphylaxis, have been reported in KANUMA-treated patients. In clinical trials, 3 of 106 (3%) patients treated with KANUMA experienced signs and symptoms consistent with anaphylaxis. These patients experienced reactions during infusion with signs and symptoms including chest discomfort, conjunctival injection, dyspnea, generalized and itchy rash, hyperemia, swelling of eyelids, rhinorrhea, severe respiratory distress, tachycardia, tachypnea, and urticaria. Anaphylaxis has occurred as early as the sixth infusion and as late as 1 year after treatment initiation.

In clinical trials, 21 of 106 (20%) KANUMA-treated patients, including 9 of 14 (64%) infants and 12 of 92 (13%) pediatric patients, 4 years and older, and adults, experienced signs and symptoms either consistent with or that may be related to a hypersensitivity reaction. Signs and symptoms of hypersensitivity reactions, occurring in two or more patients, included abdominal pain, agitation, fever, chills, diarrhea, eczema, edema, hypertension, irritability, laryngeal edema, nausea, pallor, pruritus, rash, and vomiting. The majority of reactions occurred during or within 4 hours of the completion of the infusion. Patients were not routinely pre-medicated prior to infusion of KANUMA in these clinical trials.

Due to the potential for anaphylaxis, appropriate medical support should be readily available when KANUMA is administered. If anaphylaxis occurs, immediately discontinue the infusion and initiate appropriate medical treatment. Observe patients closely during and after the infusion. Inform patients of the signs and symptoms of anaphylaxis, and instruct them to seek immediate medical care should signs and symptoms occur.

The management of hypersensitivity reactions should be based on the severity of the reaction and may include temporarily interrupting the infusion, lowering the infusion rate, and/or treatment with antihistamines, antipyretics, and/or corticosteroids. If interrupted, the infusion may be resumed at a slower rate with increases as tolerated. Pre-treatment with antipyretics and/or antihistamines may prevent subsequent reactions in those cases where symptomatic treatment was required. If a severe hypersensitivity reaction occurs, immediately discontinue the infusion and initiate appropriate medical treatment.

Consider the risks and benefits of re-administering KANUMA following a severe reaction. Monitor patients, with appropriate resuscitation measures available, if the decision is made to re-administer the product.

## 5.2 Hypersensitivity to Eggs or Egg Products

KANUMA is produced in the egg whites of genetically engineered chickens. Patients with a known history of egg allergies were excluded from the clinical trials. Consider the risks and benefits of treatment with KANUMA in patients with known systemic hypersensitivity reactions to eggs or egg products.

## 6 ADVERSE REACTIONS

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

In clinical trials, a total of 106 patients received treatment with KANUMA. The data described below reflect exposure to KANUMA in 75 patients who received KANUMA at dosages up to 3 mg/kg once weekly in clinical trials:

- Nine patients (5 males, 4 females) who had growth failure or other evidence of rapidly progressive LAL deficiency presenting within the first 6 months of life received KANUMA for up to 165 weeks (median 60 weeks) at escalating doses ranging between 0.35 mg/kg and 5 mg/kg once weekly [see *Clinical Studies (14.1)*]. The recommended initial dosage for these patients is 1 mg/kg escalating to 3 mg/kg once weekly [see *Dosage and Administration (2.1)*].
- 66 pediatric and adult patients with LAL deficiency aged 4 to 58 years (33 males, 33 females) received KANUMA 1 mg/kg every other week up to 36 weeks.

Table 2 summarizes the most common adverse reactions occurring in >30% of patients with rapidly progressive LAL deficiency presenting within the first 6 months of life receiving KANUMA.

**Table 2: Most Common Adverse Reactions\* in Patients with Rapidly Progressive LAL Deficiency Presenting within the First 6 Months of Life**

Adverse Reactions	KANUMA N=9
	n (%)
Diarrhea	6 (67)
Vomiting	6 (67)
Fever	5 (56)
Rhinitis	5 (56)
Anemia	4 (44)
Cough	3 (33)
Nasopharyngitis	3 (33)
Urticaria	3 (33)

\* Reported in more than 30% of patients receiving KANUMA

Other less common adverse reactions reported in patients with rapidly progressive disease presenting within the first 6 months of life who received KANUMA included hypotonia, decreased oxygen saturation, retching, sneezing, and tachycardia.

Table 3 summarizes the most common adverse reactions that occurred in ≥8% of pediatric and adult patients with LAL deficiency receiving KANUMA at a dosage of 1 mg/kg once every other week during the 20-week double-blind treatment period [see *Clinical Studies (14.2)*].

**Table 3: Most Common Adverse Reactions\* in Pediatric and Adult Patients with LAL Deficiency**

Adverse Reactions	KANUMA N = 36	Placebo N = 30
	n (%)	n (%)
Headache	10 (28)	6 (20)
Fever	9 (25)	7 (23)
Oropharyngeal pain	6 (17)	1 (3)
Nasopharyngitis	4 (11)	3 (10)
Asthenia	3 (8)	1 (3)
Constipation	3 (8)	1 (3)
Nausea	3 (8)	2 (7)

\* Reported in at least 8% of pediatric and adult patients receiving KANUMA and at a higher incidence than in patients receiving placebo

Other less common adverse reactions reported in pediatric and adult patients who received KANUMA included anxiety and chest discomfort.

### Hyperlipidemia

Increases in circulating LDL-cholesterol (LDL-c) and triglycerides above pre-treatment values were observed in 29 of 36 (81%) and 21 of 36 (58%) patients, respectively, at 2 and 4 weeks following initiation of KANUMA [see *Clinical Pharmacology (12.2)*]. The maximum mean percentage increase was 18% for LDL-c at Week 2 and 5% for triglycerides at Week 4.

### 6.2 Immunogenicity

As with all therapeutic proteins, there is potential for immunogenicity. Patients have developed anti-drug antibodies (ADA) to KANUMA. Immunogenicity assay results are highly dependent on the sensitivity and specificity of the assay and may be influenced by several factors such as: assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies to KANUMA with the incidence of antibodies to other products may be misleading.

#### Patients with Rapidly Progressive LAL Deficiency Presenting within the First 6 Months of Life

Seven of the 9 infants with rapidly progressive disease had at least one post-treatment ADA assessment, and 4 of these 7 (57%) patients developed ADA during treatment with KANUMA. Two of the 4 ADA-positive patients were determined to be positive for neutralizing antibodies that inhibit *in vitro* enzyme activity and cellular uptake of the enzyme. At the time of initial ADA positivity, 3 patients were receiving a dosage of 1 mg/kg once weekly and 1 patient was receiving a dosage of 3 mg/kg once weekly. Three of the 4 ADA-positive patients had ADA titers monitored from the initiation of treatment, and developed measurable ADA titers within the first 2 months of exposure. One of the 4 ADA-positive patients had persistent ADA titers. ADA titers decreased to undetectable levels in the remaining 3 patients while receiving continued treatment at a dosage of 3 mg/kg once weekly.

Hypersensitivity reactions occurred in all 4 of the ADA-positive patients, whereas they occurred in only 1 of the 3 ADA-negative patients. None of the patients discontinued treatment. In 1 patient, decreased growth velocity in a setting of neutralizing antibodies to KANUMA was observed.

#### Pediatric and Adult Patients with LAL Deficiency

Five of 35 (14%) KANUMA-treated pediatric and adult patients who completed the 20-week double-blind period of study treatment developed ADA. All patients were receiving 1 mg/kg once every other week. All 5 ADA-positive patients first developed measurable ADA titers within the first 3 months of exposure. Two of the 5 ADA-positive patients had a measurable ADA titer at only one time point. In the 3 patients with measurable ADA titers at multiple time points, ADA titers decreased to undetectable levels during continued treatment. Two patients developed *in vitro* neutralizing antibodies during the open-label extension phase after 20 weeks and 52 weeks of treatment with KANUMA, respectively. There is no clear association between the development of ADA and decreased efficacy in pediatric and adult patients treated with KANUMA.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

There are no available data on KANUMA in pregnant women to inform any drug-associated risk. Animal reproductive studies conducted with sebelipase alfa showed no evidence of embryoletality, fetotoxicity, teratogenicity, or abnormal early embryonic development at dosages up to 164 and 526 times the human dosage of 1 mg/kg every other week (based on AUC) in rats and rabbits, respectively.

The background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

#### Animal Data

Sebelipase alfa administered during the period of organogenesis to rats (on gestation days 6, 9, 12, 15 and 17) and rabbits (on gestation days 7, 10, 13, 16 and 19) at intravenous doses up to 60 and 50 mg/kg, respectively, (approximately 164 and 526 times the human AUC of 1387 ng.h/mL at 1 mg/kg dose administered once every other week, respectively) did not cause any adverse effects on embryofetal development. A pre- and postnatal development study in rats showed no evidence of adverse effects on pre- and postnatal development at intravenous doses (administered on gestation days 6, 9, 12, 15, 18, and 20 and days 4, 7, 10, 14, and 17 postpartum) of sebelipase alfa up to 60 mg/kg/day (approximately 164 times the human AUC of 1387 ng.h/mL at 1 mg/kg dose administered once every other week).

### 8.2 Lactation

#### Risk Summary

There are no data on the presence of sebelipase alfa in human milk, the effects on the breastfed infant, or the effects on milk production. It is not known if sebelipase alfa is present in animal milk. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for KANUMA and any potential adverse effects on the breastfed infant from sebelipase alfa or from the underlying maternal condition.

### 8.4 Pediatric Use

Safety and effectiveness of KANUMA have been established in pediatric patients aged 1 month and older. Clinical trials with KANUMA were conducted in 56 pediatric patients (range 1 month to <18 years old) [see *Clinical Studies (14)*].

### 8.5 Geriatric Use

Clinical trials of KANUMA did not include any patients aged 65 years old and older. It is not known whether they respond differently than younger patients.

## 11 DESCRIPTION

KANUMA (sebelipase alfa) is a recombinant human lysosomal acid lipase (rhLAL). Lysosomal acid lipase (EC 3.1.1.13) is a lysosomal glycoprotein enzyme that catalyzes the hydrolysis of cholesteryl esters to free cholesterol and fatty acids and the hydrolysis of triglycerides to glycerol and free fatty acids.

KANUMA is produced by recombinant DNA technology in the egg white of eggs laid by genetically engineered chickens. Purified sebelipase alfa is a monomeric glycoprotein containing 6 N-linked



glycosylation sites and has a molecular mass of approximately 55,000 daltons. The amino acid sequence for sebelipase alfa is the same as the amino acid sequence for human LAL. The specific activity of sebelipase alfa is 195 to 345 units/mg. One unit is the amount of enzyme activity that catalyzes the hydrolysis of 1 micromole of the synthetic substrate 4-methylumbelliferyl oleate per minute at 37°C under specified assay conditions.

KANUMA is supplied as a sterile, preservative-free, non-pyrogenic aqueous solution in single-use vials for intravenous infusion. Each vial contains sebelipase alfa 20 mg/10 mL. Each mL of solution contains sebelipase alfa (2 mg), citric acid monohydrate (1.57 mg), Human Serum Albumin (10 mg), and trisodium citrate dihydrate (13.7 mg) at pH 5.9.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

LAL deficiency is an autosomal recessive lysosomal storage disorder characterized by a genetic defect resulting in a marked decrease or loss in activity of the lysosomal acid lipase (LAL) enzyme. The primary site of action of the LAL enzyme is the lysosome, where the enzyme normally causes the breakdown of lipid particles including LDL-c. Deficient LAL enzyme activity results in progressive complications due to the lysosomal accumulation of cholesteryl esters and triglycerides in multiple organs, including the liver, spleen, intestine, and the walls of blood vessels. The resulting lipid accumulation in the liver may lead to increased liver fat content and progression of liver disease, including fibrosis and cirrhosis. Lipid accumulation in the intestinal wall leads to malabsorption and growth failure. In parallel, dyslipidemia due to impaired degradation of lysosomal lipid is common with elevated LDL-c and triglycerides and low HDL-cholesterol (HDL-c).

Sebelipase alfa binds to cell surface receptors via glycans expressed on the protein and is subsequently internalized into lysosomes. Sebelipase alfa catalyzes the lysosomal hydrolysis of cholesteryl esters and triglycerides to free cholesterol, glycerol and free fatty acids.

### 12.2 Pharmacodynamics

In clinical trials, after initiation of dosing with KANUMA, breakdown of accumulated lysosomal lipid led to initial increases in LDL-c and triglycerides within the first 2 to 4 weeks of treatment. In general, following increases in LDL-c and triglycerides, these parameters decreased to below pre-treatment values within 8 weeks of treatment with KANUMA.

In all patients with elevated alanine aminotransferase (ALT) values at baseline (82 of 84 patients in clinical trials), reductions in ALT values were observed, generally within 2 weeks after initiation of treatment with KANUMA. Treatment interruption resulted in increases in LDL-c and ALT values and decreases in HDL-c.

### 12.3 Pharmacokinetics

The pharmacokinetic profile of sebelipase alfa was nonlinear with a greater than dose-proportional increase in exposure between 1 and 3 mg/kg based on non-compartmental analysis of data from 26 adults. No accumulation was observed following once weekly or once every other week dosing. Using a population pharmacokinetic model, sebelipase alfa pharmacokinetic parameters were estimated for 65 pediatric and adult patients who received intravenous infusions of KANUMA at 1 mg/kg at Week 22 (Table 4); 24 patients were 4 to 11 years old, 23 were 12 to 17 years old, and 18 were adults. The pharmacokinetic profiles of sebelipase alfa were similar between adolescents and adults. The  $T_{max}$  and  $T_{1/2}$  were similar across all age groups.

**Table 4: Mean (SD) Pharmacokinetics Parameters at Week 22 in Pediatric and Adult Patients Receiving 1 mg/kg Once Every Other Week**

Parameter	4-11 years old	12-17 years old	≥18 years old
	N=24	N=23	N=18
AUC (ng·hr/mL)	942 (388)	1454 (699)	1861 (599)
$C_{max}$ (ng/mL)	490 (205)	784 (480)	957 (303)
$T_{max}$ (hr)	1.3 (0.6)	1.1 (0.3)	1.3 (0.6)
CL (L/hr)	31.1 (7.1)	37.4 (12.4)	38.2 (12.5)
$V_c$ (L)	3.6 (3.0)	5.4 (2.4)	5.3 (1.6)
$T_{1/2}$ (min)	5.4 (4.3)	6.6 (3.7)	6.6 (3.7)

Parameter values were estimated using a population pharmacokinetic model.

AUC = Area under the plasma concentration time curve.  $C_{max}$  = Maximum concentration.

$T_{max}$  = Time to maximum concentration. CL = Clearance.  $V_c$  = Central volume of distribution.  $T_{1/2}$  = Half-life.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals to evaluate carcinogenic potential or studies to evaluate mutagenic potential have not been performed with sebelipase alfa. Sebelipase alfa at intravenous doses up to 60 mg/kg administered twice weekly (approximately 164 times the human AUC of 1387 ng·h/mL at 1 mg/kg dose administered once every other week) was found to have no adverse effect on fertility and reproductive performance of male and female rats.

### 13.2 Animal Toxicology and/or Pharmacology

In a rat disease model of LAL deficiency that exhibits several abnormalities analogous to the human disease, sebelipase alfa administered intravenously at dosages up to 3 mg/kg once weekly showed improvements in survival, body weight gain, organ weight reduction, reduction in serum transaminases (ALT and aspartate aminotransferase [AST]), reduction in serum and hepatic lipids, and improvement in liver histopathology.

## 14 CLINICAL STUDIES

### 14.1 Patients with Rapidly Progressive LAL Deficiency Presenting within the First 6 Months of Life

A multicenter, open-label, single-arm clinical study of KANUMA was conducted in 9 infants with LAL deficiency who had growth failure or other evidence of rapidly progressive disease prior to 6 months of age. The age range at entry was 1 to 6 months. Patients received KANUMA at 0.35 mg/kg once weekly for the first 2 weeks and then 1 mg/kg once weekly. Due to suboptimal clinical response, doses in all 6 surviving patients were escalated to 3 mg/kg once weekly, between

4 and 88 weeks (median 11 weeks) after starting treatment at 1 mg/kg. In one patient, the dose was escalated to 5 mg/kg once weekly at Week 88 due to decreased growth velocity in a setting of positive neutralizing anti-drug antibodies to KANUMA. The recommended dosage for these patients is 1 mg/kg to 3 mg/kg once weekly [see *Dosage and Administration* (2.1)].

Efficacy of KANUMA was assessed by comparing the survival of 9 KANUMA-treated patients at 12 months of age with an untreated historical cohort of 21 patients with a similar age at disease presentation and clinical characteristics. Of the 9 KANUMA-treated infants, 6 patients survived beyond 12 months of age, compared to 0 of 21 patients in the historical cohort, all of whom died by 8 months of age. The median age of the 6 surviving KANUMA-treated patients was 18.1 months (range 12 to 42.2 months).

Following initiation of treatment with KANUMA 1 mg/kg once weekly, weight-for-age z-scores improved in 3 of 5 surviving patients with growth failure, and all 6 surviving patients demonstrated improvements in weight-for-age z-scores following dose escalation to 3 mg/kg once weekly.

### 14.2 Pediatric and Adult Patients with LAL Deficiency

The safety and efficacy of KANUMA were assessed in 66 pediatric and adult patients with LAL deficiency, aged 4 to 58 years (71% were less than 18 years old), in a multicenter, double-blind, placebo-controlled trial. Patients were randomized to receive KANUMA at a dosage of 1 mg/kg (n=36) or placebo (n=30) once every other week for 20 weeks in the double-blind period. Sixty-two of the 66 (94%) patients had LDL-c of 130 mg/dL or greater at study entry. The majority of patients (58%) had LDL-c above 190 mg/dL at study entry, and 24% of patients with LDL-c above 190 mg/dL remained on lipid lowering medications.

At the completion of the 20-week double-blind period of the trial, a statistically significant improvement in percent change from baseline in LDL-c was observed in the KANUMA-treated group as compared to the placebo group (mean difference and 95% C.I.: -22%, [-33%, -15%];  $p < 0.0001$ ). LDL-c of less than 130 mg/dL was achieved in 13 of 32 (41%; 95% C.I.: [24%, 58%]) KANUMA-treated patients and in only 2 of 30 (7%; 95% C.I.: [0%, 16%]) placebo-treated patients with baseline LDL-c of 130 mg/dL or greater. A statistically significant improvement in percent change from baseline at 20 weeks was also observed in the KANUMA-treated group compared to the placebo group for other parameters related to LAL deficiency, including decreases in non-HDL-c (mean difference and 95% C.I.: -21%, [-30%, -15%];  $p < 0.0001$ ) and triglycerides (mean difference and 95% C.I.: -14%, [-28%, -1%];  $p = 0.0375$ ), and increases in HDL-c (mean difference and 95% C.I.: 20%, [12%, 26%];  $p < 0.0001$ ). The effect of KANUMA on cardiovascular morbidity and mortality has not been established.

Patients treated with KANUMA had larger reductions from baseline in ALT values and liver fat content (measured by MRI), compared to patients treated with placebo. The significance of these findings as they relate to progression of liver disease in LAL deficiency has not been established.

### Open-label Extension

Pediatric and adult patients who participated in the randomized, placebo-controlled trial were eligible to continue treatment in an open-label extension. Sixty-five of 66 patients (98%) entered the open-label period in which all patients received KANUMA at a dosage of 1 mg/kg once every other week. During the open-label extension, patients treated with KANUMA for up to 36 weeks demonstrated improvements in lipid parameters, including LDL-c and HDL-c levels, and ALT.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

KANUMA 20 mg/10 mL vials are supplied as a sterile, preservative-free, nonpyrogenic solution in single-use, glass vials.

NDC 25682-007-01: 20 mg/10 mL vial

Store KANUMA under refrigeration between 2°C to 8°C (36°F to 46°F) in original carton to protect from light. Do not shake or freeze the vials.

## 17 PATIENT COUNSELING INFORMATION

### Hypersensitivity Reactions, including Anaphylaxis

Advise patients and caregivers that reactions related to administration and infusion may occur during and after KANUMA treatment, including life-threatening anaphylaxis and severe hypersensitivity reactions. Inform patients of the signs and symptoms of anaphylaxis and hypersensitivity reactions, and have them seek immediate medical care should signs and symptoms occur [see *Warnings and Precautions* (5.1)].

Manufactured by:  
Alexion Pharmaceuticals Inc.  
Cheshire, CT 06410  
US License Number: 1743  
1-888-765-4747 (phone)

KANUMA is a trademark of Alexion Pharmaceuticals Inc.